

Update of Calibration Information Used to Correct Himawari-8 AHI Sensitivity Trend

Meteorological Satellite Center  
Japan Meteorological Agency

The Japan Meteorological Agency (JMA) plans to update the calibration information used to correct the Himawari-8 sensor sensitivity trend at 07:00 UTC on 8 July 2019.

The corrected calibration parameters are included in the calibration information block in Himawari Standard Data (HSD). These are described as Nos. 12 and 13 in the #5 calibration information block for the visible and near-infrared bands (Bands 1 – 6) in HSD format<sup>1</sup>. The values of the parameters are shown in Tables 1 and 2.

JMA revised the HSD format on 25 July 2017, which includes the latest calibration coefficients with sensor sensitivity trends taken into account. Fig. 1 shows these trends for the Himawari-8 Advanced Himawari Imager (AHI-8) visible and near-infrared (VNIR) bands (i.e., Bands 1 to 6) as derived from solar diffuser observations. Degradations of approximately 0.5% per year are observed in the trends of Bands 1 to 4.

For evaluation of these trends, the sensor sensitivity correction coefficient D can be defined as

$$D = \frac{\widehat{m_{yyyy}}}{\widehat{m_{2015}}}$$

where  $\widehat{m_{yyyy}}$  is the average calibration slope for all detectors of each band as derived from four solar diffuser observations conducted on 7 and 22 May and on 7 and 22 June yyyy, while  $\widehat{m_{2015}}$  is the average calibration slope for 2015 as derived in the same way.

Multiplying the slope (no. 8 in the #5 calibration information block for HSD format) and intercept (no. 9 in the #5 calibration information block for HSD format) for 2015 by the sensor sensitivity correction coefficient D gives the corrected slope and intercept. The results are shown in Tables 1 and 2. Fig. 2 also shows the corrected sensor sensitivity trends. The degradations seen in the trends of Bands 1 to 4 in Fig. 1 are appropriately corrected.




Fig. 1 Himawari-8 sensor sensitivity trends. Time-series representations of sensor sensitivity (inverse of calibration slope) as derived from AHI-8 solar diffuser observations. Values are averaged over the detectors and normalized with the first observation made on 7 March 2015. The range of the x axis is from 7 March to 22 February.

<sup>1</sup>[https://www.data.jma.go.jp/mscweb/en/himawari89/space\\_segment/hsd\\_sample/HS\\_D\\_users\\_guide\\_en\\_v13.pdf](https://www.data.jma.go.jp/mscweb/en/himawari89/space_segment/hsd_sample/HS_D_users_guide_en_v13.pdf)




Fig. 2 Himawari-8 corrected sensor sensitivity trends. As per Fig. 1 but with correction of sensor sensitivity trends

Table 1 Slope of digital count – radiance conversion equation (Himawari-8). The latest data will be included in no. 12 of visible, near-infrared bands (Bands 1 – 6) in the #5 calibration information block for HSD format.

Band/Year	2015	2016	2017	2018	2019
B01	0.37735835	0.37920237	0.38083577	0.38225655	0.38375996
B02	0.35410388	0.35598556	0.35748863	0.35863737	0.35968951
B03	0.30549747	0.30731905	0.30913652	0.31078894	0.31231127
B04	0.18197547	0.18294331	0.18397175	0.18494062	0.18600134
B05	0.04537718	0.04536906	0.04542336	0.04540857	0.04543758
B06	0.01406841	0.01406430	0.01407068	0.01407028	0.01407496

Table 2 Intercept of digital count – radiance conversion equation (Himawari-8). The latest data will be included in no. 13 of visible, near-infrared bands (Bands 1 – 6) in the #5 calibration information block for HSD format.

Band/Year	2015	2016	2017	2018	2019
B01	-7.54716706	-7.58404731	-7.61671534	-7.64513097	-7.67519925
B02	-7.08207765	-7.11971124	-7.14977261	-7.17274746	-7.19379019
B03	-6.10994941	-6.14638096	-6.18273038	-6.21577883	-6.24622538
B04	-3.63950941	-3.65886614	-3.67943502	-3.69881245	-3.72002677
B05	-0.90754353	-0.90738115	-0.90846722	-0.90817149	-0.90875151
B06	-0.28136824	-0.28128597	-0.28141362	-0.28140566	-0.28149914

#### Documentation Change Record

Issue/revision	Date	Description
1	3 July 2017	Original edition
2	9 July 2018	Updated Fig. 1, Fig. 2, Table 1 and Table 2 to include analysis results from July 2017 to June 2018.
3	3 July 2019	Updated Fig. 1, Fig. 2, Table 1 and Table 2 to include analysis results from July 2018 to June 2019.