

# Introduction and Operation of iTacs (Basic)

Kazuto TAKEMURA

Tokyo Climate Center, Japan Meteorological Agency (TCC/JMA)

## 1. What's iTacs?

iTacs stands for “Interactive Tool for Analysis of the Climate System”. It is available on web browsers such as Internet Explorer, Firefox through Graphical User Interface (GUI) with no additional software or plug-ins. National Meteorological and Hydrological Services (NMHSs) can use it with personal IDs.




Fig.1 Schematic diagram for iTacs

iTacs is built on JMA servers and various types of dataset are saved in the system. Your client PC can access and use it via internet. The Japanese 55-year Reanalysis (JRA-55, Kobayashi et al. 2015) and outgoing longwave radiation (OLR) provided by NOAA can be used for atmospheric analysis. COBE-SST (Ishii et al. 2005) is also available in oceanographic analysis. The detailed elements available on iTacs are listed in APPENDIX.

Atmospheric analysis	JRA-55 (Kobayashi et al. 2015)	From 1958 to present
	OLR provided by NOAA	From 1979 to present
Oceanographic analysis	COBE-SST (Ishii et al. 2005)	From 1891 to present
	MOVE/MRI.COM-G2 (Toyoda et al. 2013)	From 1958 to present
Atmospheric forecast	Output of JMA's one-month prediction model	
Others	ENSO monitoring indices, CLIMAT reports, user-input data etc.	

Table.1 Available dataset on iTacs

In iTacs, various types of charts such as two-dimensional map, cross section diagram, timeseries graph can be drawn, and some types of statistical analyses such as Empirical Orthogonal Function (EOF) analysis, regression or correlation analyses can be performed. iTacs is one of the most useful tool and will help you to understand climate system.




Fig.2 Various types of charts available on iTacs


## 2. Application for using iTacs

Registered users can access iTacs at the TCC website. User ID and password are needed to use the iTacs\*. JMA permits persons at NMHSs to use the iTacs. If you are interested to use the iTacs, access the following URL for the application.

iTacs: <http://extreme.kishou.go.jp/tool/itacs-tcc2015/>

You can see “Requests for iTacs access” section on this web page. Please carefully read the conditions of use and disclaimer. If you agree to them, please applying to TCC ([tcc@met.kishou.go.jp](mailto:tcc@met.kishou.go.jp)) by e-mail completely filling the items. JMA will examine applications and, if the application is accepted, issue ID and password.

\* ID and password of the seminar participants are already issued.



iTacs:  
<http://extreme.kishou.go.jp/tool/itacs-tcc2015/>

Fig.3 Access to iTacs

## 3. Basic operations

If you input ID and password provided on the iTacs login page, you will see main display of the iTacs as shown below. The standard procedure for drawing a chart by iTacs is as follows:

- ◆ Select dataset, element, and data type.
- ◆ Set geophysical parameters: area (longitude and latitude), pressure level or depth.
- ◆ Set chronological parameters: average period (e.g., daily, monthly), period to show (e.g., year, month, day).
- ◆ Select analysis method (if needed).
- ◆ Set graphic parameters (if needed).
- ◆ Click a submit button and draw a map.




Fig.4 Main display of iTacs

### 3.1. Longitude-latitude map

As a starter, let's chart monthly sea surface temperature (SST) map in December 2015. Set parameters on "Data1" box as shown below.

- ◆ Dataset: SST (COBE-SST).
- ◆ Element: Sea Surface Data → Temperature (SST).
- ◆ Data type: HIST (historical actual observation or analysis).
- ◆ Area: ALL (90°S – 90°N, 0° – 360°E).
- ◆ Level: 1 (Surface data).
- ◆ Time unit: MONTHLY.
- ◆ Showing period: RANGE, 2015 – 12 (December 2015) for both upper and lower boxes.

Fig.5a Parameter setting on iTacs to draw SST map in December 2015

Finally, click "Analysis Data Submit" button and the image will be displayed.





Fig.5b Image of SST in December 2015 created by the setting shown in Fig.5a

You can select the following options in "Data type" pull-down menu.

- ◆ HIST: Historical actual analysis or observation data.
- ◆ NORM: Climatological normal data (averaged from 1981 to 2010).
- ◆ ANOM: Anomaly data (HIST minus NORM: difference from the climatological normal)
- ◆ ANOM\_SD: Anomaly data normalized by their standard deviations.

Select "ANOM" in "Datatype" box to draw anomalies (Fig.6a). Changing "Color Table" and "Contour Parameters" in "Graphic Options" tab as shown in Fig.6b, it becomes easier to recognize the above- and below-normal SST anomalies.



(c) Graphic options for Fig.6(b)




Fig.6 SST anomalies in December 2015 with (a) the color table “Rainbow” (default setting), (b) “Blue–Red” and (c) Graphic options of the contour setting for (b)

You can adjust zonal and meridional range by setting “Lat” and “Lon” parameters in the “Area” field. The negative values of latitude and longitude mean south latitude and west longitude, respectively (See Fig.7).




Fig.7 Example of area setting on iTacs

The following options are available in “Showing period” pull-down menu to pick up the time range to show.

- ◆ RANGE: Set the start and end points of the targeted time period.
- ◆ YEARS: Set individual years.
- ◆ INDEX: Pick up years based on a condition of SST index (e.g., NINO.3, IOBW).

### 3.2. Overlaying two data

Users can overlay two kinds of elements on the same image by using “DATA1\_DATA2” analysis method. Let’s chart and superimpose three-month mean 850-hPa stream function and its anomalies from December 2015 to February 2016 on a map. Set parameters on “Data1” and “Data2” box as shown below (See also Fig.8a).

#### ◆ Data1

- Dataset: JRA-55.
- Element: Pressure Levels →  $\psi$  (Stream Function).
- Data type: ANOM.
- Area: Lat: -40 – 40, Lon: 100 – 300 (40°S – 40°N, 100°E – 60°W).
- Level: 850 hPa.
- Time unit: MONTHLY, check “Ave” box to calculate three-month mean.
- Showing period: RANGE, 2015 – 12 for upper box and 2016 – 2 for lower box.

“Data2” box will appear after selecting “DATA1\_DATA2” in “Analysis method” pull-down menu.

#### ◆ Data2

- Data type: HIST.
- Other parameters are the same as Data1.

#### ◆ Graphic Options

- Color Table: Blue – Red.
- Set Contour Parameters for data1: interval:1, min:-5.5, max:5.5.

Fig.8a Parameter setting and graphic options to draw 850-hPa stream function and its anomalies map in DJF 2015/2016

Finally, click “Analysis Data Submit” button and the image will be displayed.




Fig.8b Image of 850-hPa stream function in DJF 2015/2016 created by the setting shown in Fig.8a

The Data1 will be mapped as shading, and Data2 is mapped as contour lines. As an exception, Data2 is mapped as shading when Data1 is mapped as the type of vector or streamline (not shown in this lecture).

### 3.3. Mapping the difference of two data

Users can calculate and map the difference of two data by using “SUBTRACT” analysis method. Let’s chart monthly SST anomaly change from October to December 2015. Set parameters on “Data1” and “Data2” box as shown below (See also Fig.9a).

#### ◆ Data1

- Dataset: SST.
- Element: Sea Surface Data → Temperature (SST).
- Data type: ANOM.
- Area: Lat: -90 – 90, Lon: -30 – 330 (90°S – 90°N, 30°W – 330°E).
- Level: 1.
- Time unit: MONTHLY.
- Showing period: RANGE, 2015 – 12 for both upper and lower boxes.

“Data2” box will be adjustable after selecting “SUBTRACT” in “Analysis method” pull-down menu.

#### ◆ Data2

- Showing period: RANGE, 2015 – 10 for both upper and lower boxes.
- Other parameters are the same as Data1.

#### ◆ Graphic Options

- Color Table: Blue – Red.
- Set Contour Parameters for data1: interval:0.2, min:-1.1, max:1.1.

In the setting above, the value of “Data1” minus “Data2” will be calculated and mapped. Click “Analysis Data Submit” button and the image will be displayed.

### Data1

Dataset	Element	Data type	Area	Level	Time unit	Showing period
SST	Sea Surface Data Temperature (SST) [t]	ANOM	ALL Lat: -90 - 90 Ave <input type="checkbox"/> Lon: -30 - 330 Ave <input type="checkbox"/>	1	MONTHLY <input type="checkbox"/> Ave <input type="checkbox"/> Year-to-year <input type="checkbox"/> Time filter	RANGE 2015 12 2015 12

Vector  SD  
Derivative:  lon  lat

Analysis method: SUBTRACT

### Data2

Dataset	Element	Data type	Area	Level	Time unit	Showing period
SST	Sea Surface Data Temperature (SST) [t]	ANOM	ALL Lat: -90 - 90 Ave <input type="checkbox"/> Lon: -30 - 330 Ave <input type="checkbox"/>	1	MONTHLY <input type="checkbox"/> Ave <input type="checkbox"/> Year-to-year <input type="checkbox"/> Time filter	RANGE 2015 10 2015 10

SD

Select parameters **Graphic Options**

### Graphic Options

Colorizing: COLOR	<input checked="" type="checkbox"/> Show Contour Labels
Drawing: SHADE	<input checked="" type="checkbox"/> Show Color Bar
Image Format: png	<input checked="" type="checkbox"/> Set Contour Parameters for data1
Font: default	interval: 0.2 min: -1.1 max: 1.1
Color Table: Blue - Red	<input type="checkbox"/> Set Vector size: [ ] [inch] value: [ ] skip: 1

Detailed Options for Image 1

Fig.9a Parameter setting and graphic options to draw monthly SST anomaly difference between October and December 2015




Fig.9b Image of SST anomaly change from October to December 2015 created by the setting shown in Fig.9a

In a similar way, users can also perform the four basic arithmetic operations of two data by using the corresponding analysis method shown in Table.2.

Analysis method	Mapped value	Usage example
ADD	Addition (“Data1” plus “Data2”)	–
SUBTRACT	Difference (“Data1” minus “Data2”)	Time difference, vertical shear.
MULTIPLY	Multiplication (“Data1” times “Data2”)	–
DIVIDE	Division (“Data1” divided by “Data2”)	Precipitation ratios (“HIST” divided by “NORM”).

Table.2 Analysis method of the four basic arithmetic operations on iTacs

#### 4. Advanced operations

Users can create various types of image such as line graph and cross section diagram (Fig.10). These operations are useful to see the variability or spatial structure simply.




Fig.10 Example of time-series graphs available on iTacs

#### 4.1. Area-averaged time series

Daily, monthly and inter-annual time-series are available by adjusting “Area”, “Level”, “Time unit” and “Showing period”. As an example, let’s chart daily time-series of area-averaged OLR anomalies. Set parameters on “Data1” box as shown below (See also Fig.11a).

- ◆ Dataset: SAT (OLR is available by selecting “SAT” in this box).
- ◆ Element: OLR [W/m<sup>2</sup>].
- ◆ Data type: ANOM.
- ◆ Area: Lat: -10 – 10, Lon: 90 – 150 (over and around the Maritime Continent).
  - Check “Ave” box in both “Lat” and “Lon” fields to calculate area-averaged value.
- ◆ Level: 1 (OLR is surface data).
- ◆ Time unit: DAILY.
- ◆ Showing period: RANGE,
  - Upper box: 2016 – 5 – 1 (from 1 May 2016),
  - Lower box: 2016 – 8 – 31 (to 31 August 2016).

### Data 1

Dataset	Element	Data type	Area	Level	Time unit	Showing period
SAT	OLR [W/m <sup>2</sup> ]	ANOM	ALL	1	DAILY	RANGE
			Lat: -10 - 10 Ave <input checked="" type="checkbox"/>			2016 5 1
			Lon: 90 - 150 Ave <input checked="" type="checkbox"/>			2016 8 31
<input type="checkbox"/> Vector <input type="checkbox"/> SD Derivative: <input type="checkbox"/> lon <input type="checkbox"/> lat					<input type="checkbox"/> Ave <input type="checkbox"/> Year-to-year <input type="checkbox"/> Time filter	

Fig.11a Parameter setting on iTacs to draw time-series of OLR anomalies from 1 May to 31 August 2016  
 Finally, click “Analysis Data Submit” button and the image will be displayed.






Fig.11b Daily time-series of OLR anomalies created by the setting shown in Fig.11a

OLR is one of the most important indices of tropical convective activities. It can be assumed that lower values of OLR indicate more enhanced convective activities, except for the mid-latitudes in winter season and the high-latitudes. In Fig.11b, negative (positive) anomalies indicate that the convective activities are stronger (weaker) than normal.

Selecting “MONTHLY” in “Time unit” box, users can draw monthly time-series.

Next, let’s chart inter-annual time-series of 200-hPa velocity potential anomalies. Set parameters on “Data1” box as shown below (See also Fig.12a).

- ◆ Dataset: JRA-55.
- ◆ Element: Pressure Levels →  $\chi$  (Velocity Potential).
- ◆ Data type: ANOM.
- ◆ Area: Lat: -10 – 10, Lon: 90 – 150 (over and around the Maritime Continent).
  - Check “Ave” box in both “Lat” and “Lon” fields to calculate area-averaged value.
- ◆ Level: 200hPa for both upper and lower boxes.
- ◆ Time unit: MONTHLY.
  - Check “Year-to-year” box to calculate inter-annual variabilities.
- ◆ Showing period: RANGE, Year: 1958 – 2016, Month: 5 – 8 (MJJA mean from 1958 to 2016).

**Data 1**

Dataset	Element	Data type	Area	Level	Time unit	Showing period
JRA-55	Pressure Levels $\chi$ (Velocity Potential)	ANOM	ALL Lat: -10 - 10 Ave <input checked="" type="checkbox"/> Lon: 90 - 150 Ave <input checked="" type="checkbox"/>	200hPa 200hPa	MONTHLY <input type="checkbox"/> Ave <input checked="" type="checkbox"/> Year-to-year <input type="checkbox"/> Time filter	RANGE 1958 - 2016 5 - 8

Vector  SD  
Derivative:  lon  lat

Fig.12a Parameter setting on iTacs to draw inter-annual time-series of four-month (May – August) averaged 200-hPa velocity potential anomalies over and around the Maritime Continent from 1958 to 2016

Finally, click “Analysis Data Submit” button and the image will be displayed.




Fig.12b Inter-annual time-series of 200-hPa velocity potential anomalies created by the setting shown in Fig.12a

#### 4.2. Vertical and latitude/longitude profile

As with the time-series graph, users can make vertical/horizontal profiles graph by using spatial average functions. Selecting two different levels in “Level” box, users can make the vertical profile of area-averaged elements as shown in Fig.13.

- ◆ Area: Check “Ave” in both “Lat” and “Lon” boxes to calculate area-averaged value.
- ◆ Level: 1000 hPa – 1 hPa.

Checking “Logarithmic Coordinates” in the Graphic Options is recommended in vertical profile graph for pressure coordinate




Fig.13 Example of parameter setting and the figure to draw a vertical profile of height anomalies

Checking either “Ave” boxes, users can make the latitude or longitude profile of a specific level. Fig.14 shows an example of the parameter setting for a longitude profile of meridional mean values.

- ◆ Area: Check “Ave” box in “Lat” field to calculate zonal averaged values.
- ◆ Time unit, Showing period: Select a specific time or time mean.

Area		Level		Time unit		Showing period	
ASIA		500hPa		MONTHLY		RANGE	
Lat: 25	- 35	Ave <input checked="" type="checkbox"/>	500hPa	<input checked="" type="checkbox"/> Ave	<input type="checkbox"/> Year-to-year	2015	6
Lon: 90	- 180	Ave <input type="checkbox"/>		<input type="checkbox"/> Time filter		2015	7




Fig.14 Example of parameter setting on and the figure to draw a longitude profile of 500-hPa height anomalies

### 4.3. Cross section diagram

Cross section diagram is also useful to see the variability or spatial structure of atmospheric or oceanographic characteristics.




Fig.15 Example of cross section diagrams available on iTacs

Selecting two different levels in “Level” box, users can make the vertical profile of area-averaged elements as shown in Fig.16.

- ◆ Area: Check “Ave” box in “Lon” fields to calculate area-averaged value.
- ◆ Level: 1000 hPa – 100 hPa.
- ◆ Time unit, Showing period: Select a specific time or time mean.

Checking “Logarithmic Coordinates” in the Graphic Options is recommended in vertical profile graph.

Area		Level	Time unit	Showing period	
ALL		1000hPa	MONTHLY	RANGE	
Lat: -60	- 60	100hPa	<input checked="" type="checkbox"/> Ave	2016	6
Lon: 90	- 150		<input type="checkbox"/> Year-to-year	2016	8
			<input type="checkbox"/> Time filter		

### Graphic Options

Colorizing: COLOR	<input checked="" type="checkbox"/> Show Contour Labels	<input type="checkbox"/> Polar Stereographic: Nort
Drawing: SHADE	<input checked="" type="checkbox"/> Show Color Bar	<input checked="" type="checkbox"/> Logarithmic Coordinates
Image Format: png	<input type="checkbox"/> Set Contour Parameters for data1	<input type="checkbox"/> Reverse the Axes
Font: default	interval: min: max:	<input type="checkbox"/> Flip the X-axis
Color Table: Rainbow	<input type="checkbox"/> Set Vector size: [inch] value: skip: 1	<input type="checkbox"/> No Caption




Fig.16 Example of parameter setting and the figure to draw a vertical cross section of normal pressure vertical velocity

Let's chart a longitude-time cross section of meridional mean 200-hPa velocity potential anomalies. Set parameters on "Data1" box as shown below (See also Fig.17a).

- ◆ Dataset: JRA-55.
- ◆ Element: Pressure Levels →  $\chi$  (Velocity Potential).
- ◆ Data type: ANOM.
- ◆ Area: Lat: -5 – 5, Lon: 0 – 360.
  - Check "Ave" box in "Lat" field to calculate meridional mean from 5°S – 5°N.
- ◆ Level: 200hPa for both upper and lower boxes.
- ◆ Time unit: DAILY.
- ◆ Showing period: RANGE,
  - Upper box: 2015 – 5 – 1 (from 1 May 2015),
  - Lower box: 2015 – 10 – 31 (to 31 October 2015).
- ◆ Graphic Options
  - Color Table: Blue – Red.
  - Set Contour Parameters for data1: interval:3, min:-15, max:15.

### Data1

Dataset	Element	Data type	Area	Level	Time unit	Showing period
JRA-55	Pressure Levels $\chi$ (Velocity Potential)	ANOM	ALL Lat: -5 - 5 Ave <input checked="" type="checkbox"/> Lon: 0 - 360 Ave <input type="checkbox"/>	200hPa 200hPa	DAILY <input type="checkbox"/> Ave <input type="checkbox"/> Year-to-year <input type="checkbox"/> Time filter	RANGE 2015 5 1 2015 10 31

Vector  SD  
Derivative:  lon  lat

### Graphic Options

Colorizing: COLOR	<input checked="" type="checkbox"/> Show Contour Labels
Drawing: SHADE	<input checked="" type="checkbox"/> Show Color Bar
Image Format: png	<input checked="" type="checkbox"/> Set Contour Parameters for data1
Font: default	interval: 3 min: -15 max: 15
Color Table: Blue - Red	<input type="checkbox"/> Set Vector size: [inch] value: skip: 1

Fig.17a Parameter setting and graphic options to draw a longitude-time cross section of 5°S – 5°N mean 200-hPa velocity potential anomalies during the period from 1 May to 31 October 2015

Finally, click "Analysis Data Submit" button and the image will be displayed.




Fig.17b Longitude-time cross section diagram (it is a so-called “Hovmöller diagram”) created by the setting shown in Fig.17a

#### 4.4. Time filter

Time filtering is used to create a time-series or time cross section images. It emphasizes climatological variability because it can remove high frequency variations. There are two types of the time filter in iTacs as shown below:

- ◆ Running mean: Smooth the original data simply.
- ◆ Lanczos filter: Pick up the given period component and mean them based on Duchon (1979).




Fig.18 Two types of time filtering function on iTacs

Using “Running mean” function, users can make smoothed time-series graphs as shown in Fig.19a.

- ◆ Time unit: DAILY.
  - Check “Time filter” and select “Running mean”.
  - Input “5” in mean period (i.e. 5-day running mean).
- ◆ Other parameters are the same as Fig.11a.




Fig.19a As for Fig.11a, but parameter setting for 5-day running mean




Fig.19b As for Fig.11b, but smoothed by 5-day running mean created by the setting shown in Fig.19a

Smoothed time cross section diagrams are also available by using “Running mean” function as shown in Fig.20a.

- ◆ Time unit: DAILY.
  - Check “Time filter” and select “Running mean”.
  - Input “5” in mean period (i.e. 5-day running mean).
- ◆ Other parameters are the same as Fig.17a.

### Data1

Dataset	Element	Data type	Area	Level	Time unit	Showing period
JRA-55	Pressure Levels χ (Velocity Potential)	ANOM	ALL Lat: -5 - 5 Ave <input checked="" type="checkbox"/> Lon: 0 - 360 Ave <input type="checkbox"/>	200hPa 200hPa	DAILY <input type="checkbox"/> Ave <input type="checkbox"/> Year-to-year <input checked="" type="checkbox"/> Time filter Running mean	RANGE 2015 5 1 2015 10 31
<input type="checkbox"/> Vector <input type="checkbox"/> SD Derivative: <input type="checkbox"/> lon <input type="checkbox"/> lat						mean period 5

### Graphic Options

Colorizing: COLOR	<input checked="" type="checkbox"/> Show Contour Labels
Drawing: SHADE	<input checked="" type="checkbox"/> Show Color Bar
Image Format: png	<input checked="" type="checkbox"/> Set Contour Parameters for data1
Font: default	interval: 3 min: -15 max: 15
Color Table: Blue - Red	<input type="checkbox"/> Set Vector size: [inch] value: skip: 1

Fig.20a Parameter setting and graphic options to draw a longitude-time cross section of 5°S – 5°N mean 200-hPa velocity potential anomalies during the period from 1 May to 31 October 2015




Fig.20b As for Fig.17b, but smoothed by 5-day running mean created by the setting shown in Fig.20a





# Operation of iTacs (Advanced)

Yusuke URABE

Tokyo Climate Center, Japan Meteorological Agency (TCC/JMA)

## 5. Statistical Analysis

Some types of statistical analyses are available on iTacs.

- ◆ Regression/correlation analysis.
- ◆ Composite analysis.
- ◆ Single/multi EOF, Singular Value Decomposition (SVD) analysis.
- ◆ Fast Fourier Transform (FFT) analysis.
- ◆ Wavelet analysis

These methods can be powerful tools to consider and understand climate system. However, it should be noted that the results produced by them don't always mean the existence of physical system or structures in targeted data, because it just indicates mathematically calculated value and consideration based on physics isn't included.

In this text, regression/correlation analysis and composite analysis are described. These methods are frequently used to analyze the relations between global/regional climate variability and primary modes such as ENSO.

### 5.1. Regression and Correlation analysis

Regression and correlation analysis are often used to examine the circulation pattern related to the focused one-dimensional time-series. Correlation coefficient means the degree of the correlation, and the regression coefficient means the gradient of the regression line.

As an example, let's chart regression coefficient of sea level pressure (SLP) anomaly onto NINO.3 SST index for boreal winter (December to February) mean from 1958/1959 to 2010/2011. Set parameters on "Data1" box as shown below.

- ◆ Data1
  - Dataset: JRA-55.
  - Element: Surface → SLP (Sea Level Pressure).
  - Data type: ANOM.
  - Area: Lat: -90 – 90, Lon: 0 – 360.
  - Level: 1.
  - Time unit: MONTHLY, Check "Year-to-year" box in regression or correlation analysis.
  - Showing period: RANGE, 1958 – 12 for upper box and 2010 – 2 for lower box.

Be careful to parameters in "Showing period". Although the end of the range may appear to be February 2010, this setting means December to February mean from 1958/1959 (December 1958 to February 1959) to 2010/2011 (December 2010 to February 2011). Consider the setting for year and month separately.

"Data2" box will appear after selecting "REGRESSION\_COEFFICIENT" in "Analysis method" pull-down menu.

- ◆ Data2
  - Dataset: INDEX.
  - Element: NINO.3.
  - Data type: ANOM.

- Time unit: MONTHLY, Check “Year-to-year” box in regression or correlation analysis.
- Lag: 0 YEAR. (In this example, simultaneous regression is calculated)
- Significance: 95% two side. (significance test using student’s t-test)

◆ **Graphic Options**

- Drawing: CONTOUR
- Color Table: Blue – Red.
- Set Contour Parameters for data1: interval:0.4, min:-2.2, max:2.2.

### Data1

Dataset	Element	Data type	Area	Level	Time unit	Showing period
JRA-55	Surface SLP (Sea Level Press)	ANOM	ALL Lat: -90 - 90 Ave <input type="checkbox"/> Lon: 0 - 360 Ave <input type="checkbox"/>	1	MONTHLY <input type="checkbox"/> Ave <input checked="" type="checkbox"/> Year-to-year <input type="checkbox"/> Time filter	RANGE 1958 - 2010 12 - 2

Vector  SD  
Derivative:  lon  lat

Analysis method: REGRESSION\_COEFFICIENT

---

### Data2

Dataset	Element	Data type	Time unit	Lag	Significance
INDEX	NINO.3 <input type="checkbox"/> SD	ANOM	MONTHLY <input type="checkbox"/> Ave <input checked="" type="checkbox"/> Year-to-year <input type="checkbox"/> Time filter	0 YEAR	95%(two side)

### Graphic Options

Colorizing: COLOR	<input checked="" type="checkbox"/> Show Contour Labels
Drawing: CONTOUR	<input checked="" type="checkbox"/> Show Color Bar
Image Format: png	<input checked="" type="checkbox"/> Set Contour Parameters for data1
Font: default	interval: 0.4 min: -2.2 max: 2.2
Color Table: Blue - Red	<input type="checkbox"/> Set Vector size: [ ] [inch] value: [ ] skip: 1

Fig.21a Parameter setting and graphic options to draw regression coefficient and significance of SLP onto NINO.3 index for DJF 1958/1959 to 2010/2011

Finally, click “Analysis Data Submit” button and the image will be displayed.




Fig.21b Image of regression coefficient and significance of SLP onto NINO.3 index for DJF 1958/1959 to 2010/2011 created by the setting shown in Fig.21a

The regression coefficient will be mapped as contour, and the area where regression coefficient is significant at a

given (in this case, 95%) confidence level is marked as gray shading.

Correlation analysis can be done by the same process with analysis method “CORRELATION\_COEFFICIENT”.

## 5.2. Composite analysis

Users can calculate and map the composite data by using “COMPOSITE” analysis method. Let’s chart composite of temperature anomaly at 850hPa in January for years when SST anomaly in NINO.3 region was larger than +0.5. In this time, target period is from 1958 to 2010. Set parameters on “Data1” and “Data2” box as shown below (See also Fig.9a).

### ◆ Data1

- Dataset: JRA-55.
- Element: Pressure Levels → T (Temperature).
- Data type: ANOM.
- Area: Lat: -90 – 90, Lon: 0 – 360 (90°S – 90°N, 0° – 360°E).
- Level: 850hPa.
- Time unit: MONTHLY, Check “Year-to-year” box.
- Showing period: RANGE, 1958 – 1 for upper box and 2010 – 1 for lower box.

“Data2” box will appear after selecting “COMPOSITE” in “Analysis method” pull-down menu.

### ◆ Data2

- Dataset: SST
- Element: Sea Surface Data → Temperature (SST)
- Data type: ANOM. > 0.5
- Area: Lat: -5 – 5, Lon: 210 – 270 (NINO.3 region).
  - ✧ Check “Ave” box in both “Lat” and “Lon” fields to calculate area-averaged value.
- Level: 1
- Time unit: MONTHLY, Check “Year-to-year” box.

### ◆ Graphic Options

- Drawing: SHADE
- Color Table: Blue – Red.
- Set Contour Parameters for data1: interval:0.4, min:-2.2, max:2.2.

Finally, click “Analysis Data Submit” button and the image will be displayed.

## Data1

Dataset	Element	Data type	Area	Level	Time unit	Showing period
JRA-55	Pressure Levels T (Temperature) [C.I]	ANOM	ALL Lat: -90 - 90 Ave <input type="checkbox"/> Lon: 0 - 360 Ave <input type="checkbox"/>	850hPa	MONTHLY <input type="checkbox"/> Ave <input checked="" type="checkbox"/> Year-to-year <input type="checkbox"/> Time filter	RANGE 1958 - 2010 1 - 1
<input type="checkbox"/> Vector <input type="checkbox"/> SD Derivative: <input type="checkbox"/> lon <input type="checkbox"/> lat Analysis method: COMPOSITE						

## Data2

Dataset	Element	Data type	Area	Level	Time unit
SST	Sea Surface Data Temperature (SST) [t] <input type="checkbox"/> SD	ANOM > 0.5	ALL Lat: -5 - 5 Ave <input checked="" type="checkbox"/> Lon: 210 - 270 Ave <input checked="" type="checkbox"/>	1	MONTHLY <input type="checkbox"/> Ave <input checked="" type="checkbox"/> Year-to-year <input type="checkbox"/> Time filter

## Graphic Options

Colorizing: COLOR	<input checked="" type="checkbox"/> Show Contour Labels
Drawing: SHADE	<input checked="" type="checkbox"/> Show Color Bar
Image Format: png	<input checked="" type="checkbox"/> Set Contour Parameters for data1
Font: default	interval: 0.4 min: -2.2 max: 2.2
Color Table: Blue - Red	<input type="checkbox"/> Set Vector size: [ ] [inch] value: [ ] skip: 1

Detailed Options for Image 1

Fig.22a Parameter setting and graphic options to draw composite of temperature anomaly in January for years when SST anomaly in NINO.3 region was larger than +0.5.




Fig.22b Image of temperature anomaly composite at 850hPa in January created by the setting shown in

Fig.22a

If the years are already listed, you can get composite map using another settings. As an example, let's chart stream function anomaly at 850hPa in July averaged for years when El Niño events occurred in the previous winter.

### ◆ Data1

- Dataset: JRA-55.
- Element: Pressure Levels →  $\psi$  (Stream Function).
- Data type: ANOM.
- Area: Lat: -90 – 90, Lon: 0 – 360 (90°S – 90°N, 0° – 360°E).
- Level: 850hPa.
- Time unit: MONTHLY, Check “Ave” and “Year-to-year” box.
- Showing period: YEARS. Input following years in each small box or in large box with comma-separated or space-separated format.

- ✧ 1964,1966,1969,1970,1973,1977,1983,1987,1988,1992,1998,2003,2010
- ✧ Select 7 for both left and right pull-down boxes.

◆ Graphic Options

- Drawing: SHADE
- Color Table: Blue – Red.
- Set Contour Parameters for data1: interval:0.4, min:-2.2, max:2.2.

Finally, click “Analysis Data Submit” button and the image will be displayed.

**Data1**

Dataset	Element	Data type	Area	Level	Time unit	Showing period
JRA-55	Pressure Levels ψ (Stream Function)	ANOM	ALL Lat: -90 - 90 Ave <input type="checkbox"/> Lon: 0 - 360 Ave <input type="checkbox"/>	850hPa	MONTHLY <input checked="" type="checkbox"/> Ave <input checked="" type="checkbox"/> Year-to-year <input type="checkbox"/> Time filter	YEARS input years directly (comma-separated or space-separated) 1964,1966,1969,1970,1973,1977,1983,1987,1988,1992,1998,2003,2010 7 - 7

Vector  SD  
Derivative:  lon  lat

**Graphic Options**

Colorizing: COLOR	<input checked="" type="checkbox"/> Show Contour Labels
Drawing: SHADE	<input checked="" type="checkbox"/> Show Color Bar
Image Format: png	<input checked="" type="checkbox"/> Set Contour Parameters for data1
Font: default	interval: 0.4 min: -2.2 max: 2.2
Color Table: Blue - Red	<input type="checkbox"/> Set Vector size: [ ] [inch] value: [ ] skip: 1

Detailed Options for Image 1

Fig.23a Parameter setting and graphic options to draw composite of stream function anomaly at 850hPa in July for “post-EI Niño” years.




Fig.23b Image of stream function anomaly composite at 850hPa in July for post-EI Niño years created by the setting shown in Fig.23a

If you want to draw composite map using anomaly relative to default (1981-2010 mean) climatology, significance of the composite anomaly can be drawn using “SIGNIFICANCE\_TEST” analysis.

◆ Data1

- Time unit: MONTHLY, Check “Year-to-year” box.
- Other parameters are the same as previous example.

“Data2” box will appear after selecting “SIGNIFICANCE\_TEST” in “Analysis method” pull-down menu.

◆ Data2

- Showing period: RANGE, 1981 – 7 for upper box and 2010 – 7 for lower box.
- Significance: 95% (two side).
- Other parameters are the same as Data1.

◆ Graphic Options

- Drawing: CONTOUR
- Color Table: Blue – Red.
- Set Contour Parameters for data1: interval:0.4, min:-2.2, max:2.2.

Finally, click “Analysis Data Submit” button and the image will be displayed.

The image shows a web-based parameter configuration interface. It is divided into three main sections: Data1, Data2, and Graphic Options.

**Data1 Section:**

- Dataset:** JRA-55
- Element:** Pressure Levels,  $\psi$  (Stream Function)
- Data type:** ANOM
- Area:** ALL, Lat: -90 to 90, Lon: 0 to 360
- Level:** 850hPa
- Time unit:** MONTHLY,  Ave,  Year-to-year,  Time filter
- Showing period:** YEARS, with a list of years (1964, 1966, 1969, 1970, 1973, 1977, 1983, 1987, 1988, 1992, 1998, 2003, 2010) and a range of 7 years.
- Analysis method:** SIGNIFICANCE\_TEST

**Data2 Section:**

- Dataset:** JRA-55
- Element:** Pressure Levels,  $\psi$  (Stream Function)
- Data type:** ANOM
- Level:** 850hPa
- Time unit:** MONTHLY,  Ave,  Year-to-year,  Time filter
- Showing period:** RANGE, 1981 - 2010, 7 - 7, remove data1 period
- Significance:** 95%(two side)

**Graphic Options Section:**

- Colorizing:** COLOR
- Drawing:** CONTOUR
- Image Format:** png
- Font:** default
- Color Table:** Blue - Red
- Show Contour Labels
- Show Color Bar
- Set Contour Parameters for data1
- Contour Parameters:** interval: 0.4, min: -2.2, max: 2.2
- Set Vector size: [ ] [inch] value: [ ] skip: 1
- Detailed Options for Image 1

Fig.24a Parameter setting and graphic options to draw composite of stream function anomaly at 850hPa in July for “post-El Niño” years.




Fig.24b Image of stream function anomaly at 850hPa composite for post-El Niño July with 95% confidence level marked by gray shading created by the setting shown in Fig.24a

CAUTION : In “SIGNIFICANCE\_TEST” analysis, drawn map is composite of “DATA1”, not “difference between DATA1 and DATA2”. Therefore, if you want to draw composite map of deviation relative to reference value except default climatology (1981-2010 average), this method cannot be used. For example, if you change the period of DATA2 to “1971-7 to 2000-7” in the above example, the contour would indicate the composite of anomaly (relative to 1981-2010 climatology) and gray shading would indicate the significance of deviation relative to 1971-2000 average, and the figure might be misleading.

## 6. Other Functions

### 6.1. Data download

Users can download data as a zip compressed file with a plain text and GrADS format (control file and data file) to create a map. If you need to download data, please click “Download text zip file” or “Download data (ctl file and 4-byte-binary data)” at the bottom of the iTacs window.




Fig.25 Screen image of banners for download data from iTacs.

### 6.2. Using user input data

Time series data made by individual users can be used in a dataset name “USER\_INPUT”. Using this function, a correlation or regression coefficient map between single station data or user’s original index and another dataset like JRA-55 can be created, for example.

There are two ways to set data, which can be selected in element name.

- UPLOAD\_TXT : Data are given by an uploaded text file.
- INPUT\_DATA : Data are directly input to the box.

Input data must be consists of one element, separated by comma and given by specified format as “year, month, day, value”. In case of monthly data, “day” is always given as “1”. Similarly, in case of annual data, “month” and “day” are always given as “1”.

Sentences beginning with “#” have special meanings as below. Other sentences beginning with “#” is taken as comments and not used in analysis.

- “#undef =” : Definition of missing data (default is -9999). For example, “#undef=-9999”
- “#element =” : Data name used to save them on the server. For example, “#element=Daily Ts”






Fig.26 Screen image when a user inputs monthly time series data by "INPUT\_DATA".

## References

- ✓ Duchon, C. E., 1979: Lanczos Filtering in One and Two Dimensions, *J. Applied Met.*, **18**, 1016-1022.
- ✓ Ishii, M., A. Shouji, S. Sugimoto, and T. Matsumoto, 2005: Objective Analyses of Sea-Surface Temperature and Marine Meteorological Variables for the 20th Century using ICOADS and the Kobe Collection. *Int. J. Climatol.*, **25**, 865-879.
- ✓ Kobayashi, S., Y. Ota, Y. Harada, A. Ebita, M. Moriya, H. Onoda, K. Onogi, H. Kamahori, C. Kobayashi, H. Endo, K. Miyaoka, and K. Takahashi, 2015: The JRA-55 Reanalysis: General Specifications and Basic Characteristics. *J. Meteorol. Soc. Japan*, **93**, 5-48.
- ✓ Toyoda, T., Y. Fujii, T. Yasuda, N. Usui, T. Iwao, T. Kuragano, and M. Kamachi, 2013: Improved Analysis of Seasonal-Interannual Fields Using a Global Ocean Data Assimilation System, *Theoretical and Applied Mechanics Japan*, **61**, 31-48.

APPENDIX. Dataset and elements available on iTacs

Table.A1 List of analysis dataset and its elements available on iTacs

Dataset	Element	Unit		
JRA-55	Pressure Levels	$\chi$ (Velocity potential)	$10^6 \text{ m}^2/\text{s}$	
		Div (Relative divergence)	1/s	
		$\theta_e$ (Equivalent potential temperature)	K	
		$\omega$ (Pressure vertical velocity)	Pa/s	
		$\psi$ (Stream function)	$10^6 \text{ m}^2/\text{s}$	
		$\theta$ (Potential temperature)	K	
		q (Specific humidity)	kg/kg	
		T (Temperature)	$^{\circ}\text{C}$	
		T-Td (Dew point depression)	$^{\circ}\text{C}$	
		U (Zonal wind)	m/s	
		V (Meridional wind)	m/s	
		Udiv (Zonal divergence wind)	m/s	
		Vdiv (Meridional divergence wind)	m/s	
		$\zeta$ (Relative vorticity)	1/s	
		$\gamma$ (Geopotential height)	m	
	Flux	Wvf-x (Zonal water vapor flux)	kg/kg m/s	
		Wvf-y (Meridional water vapor flux)	kg/kg m/s	
	Surface	SLP (Sea level pressure)	hPa	
		Ps (Surface pressure)	hPa	
		qs (Surface specific humidity)	kg/kg	
		Ts (Surface temperature)	$^{\circ}\text{C}$	
		T-Td (Surface 2-m dew point depression)	$^{\circ}\text{C}$	
		Us (Surface zonal wind)	m/s	
		Vs (Surface meridional wind)	m/s	
		Tprat (Surface total precipitation)	mm/day	
		Latent heat flux (positive: upward)	$\text{W}/\text{m}^2$	
		Sensible heat flux (positive: upward)	$\text{W}/\text{m}^2$	
		Solar radiation flux (positive: upward)	$\text{W}/\text{m}^2$	
		Longwave radiation flux (positive: upward)	$\text{W}/\text{m}^2$	
		Net heat and radiation flux (positive: upward)	$\text{W}/\text{m}^2$	
	Isentropic Levels	Potential vorticity	PVU ( $10^{-6} \text{ K m}^2 \text{ kg}^{-1} \text{ s}^{-1}$ )	
	SST	Sea Surface Data	Temperature (SST)	$^{\circ}\text{C}$
	SAT	OLR		$\text{W}/\text{m}^2$
INDEX	NINO.1+2		$^{\circ}\text{C}$	
	NINO.3		$^{\circ}\text{C}$	
	NINO.3.4		$^{\circ}\text{C}$	
	NINO.4		$^{\circ}\text{C}$	
	NINO.WEST		$^{\circ}\text{C}$	
	IOBW		$^{\circ}\text{C}$	

Table.A2 List of forecast dataset and its elements available on iTacs

Dataset	Element	Unit	
1MONTH_ENS_MEAN	Pressure Levels	$\chi$ (Velocity potential)	$10^6 \text{ m}^2/\text{s}$
		$\psi$ (Stream function)	$10^6 \text{ m}^2/\text{s}$
		T (Temperature)	$^{\circ}\text{C}$
		T-Td (Dew point depression)	$^{\circ}\text{C}$
		U (Zonal wind)	m/s
		V (Meridional wind)	m/s
		$\gamma$ (Geopotential height)	m
	Surface	SLP (Sea level pressure)	hPa
		Ts (Surface temperature)	$^{\circ}\text{C}$
		Rain (Daily Precipitation)	mm/day
1MONTH_HIND	Pressure Levels	Velocity potential	$10^6 \text{ m}^2/\text{s}$
		Stream function	$10^6 \text{ m}^2/\text{s}$
		Geopotential height	m
		Relative humidity	%
		Temperature	$^{\circ}\text{C}$
		T-Td (Dew point depression)	$^{\circ}\text{C}$
		Zonal wind	m/s
		Meridional wind	m/s
	Surface	Surface temperature	$^{\circ}\text{C}$
		Surface pressure	hPa
		Rain	mm/day

